NEANSC WORKING PARTY ON INTERNATIONAL EVALUATION COOPERATION SUBGROUP 12: NUCLEAR MODEL VALIDATION

REPORT FROM THE 1-ST MEETING
OF SUBGROUP 12
ON
NUCLEAR MODEL VALIDATION
Gatlinburg May 8, 1994

G.Reffo May 30, 199**%**

Doc. s12. 5

Disribution list:

M. Salvatores (Chairman of the International Evaluation Cooperation NEANSC working party)

P. Oblozinsky (Monitor of subgroup 12 on Nuclear Model Vali-

dation)

A. Koning Coordinator subgroup 13

N. Kishida Coordinator subgroup 13

Y. Kikuchi Monitor subgroup 13

C. Nordborg NEA Secretariat

Participants According to list (see)

NEANSC WORKING PARTY ON INTERNATIONAL EVALUATION COOPERATION

SUBGROUP 12: NUCLEAR MODEL VALIDATION

REPORT FROM THE 1-ST MEETING
OF SUBGROUP 12
ON
NUCLEAR MODEL VALIDATION
Gatlinburg May 8, 1994

G.Reffo May 30, 1993

Introduction

At the 5-th meeting of the NEANSC working party on International Evaluation Cooperation (IEC), held at Aix-en-Provence, France, 16-th and 17-th June 1993, new subgroup activities have been established, one of which is subgroup 12 on NUCLEAR MODEL VALIDATION (NMV) (Monitor: C. Dunford, subsequently replaced by PODIozinsky; Coordinator: G. Reffo).

According to the intentions of the Iternational Evaluation Working Party [as outlined in the document NEA/NSC/WPEC(93)2, IEC-80], subgroup 12, will:

- coordinate the collection of nuclear models,
- ***
 indicate the range of applicability of each,
- produce a well-tested transportable system of computer programs to be used by the nuclear data community,
- document all the mentioned activities,
- ***
 cooperate closely with the IAEA coordinated development of a data base of nuclear model parameters.

- 1. NEED FOR NUCLEAR MODEL VALIDATION AND INTERNATIONAL COOPERATION FRAMEWORK.
- 1.1 Hystorical background and the present situation.

Low and intermediate energy nuclear physics has become a mature science and is well oriented for applications. Scientific developments in the field of both pure and applied physics are demanding nuclear data of higher and higher quality in an increasing range of masses, of reaction-types and of energies.

In the past, the bulk of nuclear data needed for technological applications was mainly provided by experimental measurement programmes. With increasing computer capabilities and availability, theoretical developments of nuclear physics have been greatly facilitated so that we start having available nuclear models and codes, as well as nuclear modelling methodologies, allowing the production of reliable nuclear data at a cost uncomparably lower than measurements.

The trend of declining funding for low and intermediate energy nuclear physics has particularly affected measurements activities, and there does not seem to be any prospect of change in the forseeable future. It will never be possible, however, to rely completely, solely, on theoretical model predictions, this is why one has to strongly support some experimental activities. Even a few measurements, in fact, could be sufficient to benchmark model assumptions, by comparison of model predictions with measurements performed in particularly critical cases, suitably selected.

In such a scenario one has to develop the capability to respond to the nuclear data demand, to a large extent, by use of theoretical and computational tools.

1.2. Necessity for a strong international cooperative effort.

Actually nuclear models and codes have been developed, which frequently rely on different physics, so that we are now faced with the necessity of clarifications. These may be achieved by means of a critical analysis and intercomparison of model results and particularly of model assumptions, as well as of the way they are used and coded.

This is a very necessary and important task which demands a strong international cooperative effort, as addressed by the IEC working party.

Indeed such an activity has already been initiated by means of extensive code intercomparisons, which will be a very useful starting point for the NMV activities.

1.3. The link to the IAEA coordinated development of a REFERENCE INPUT PARAMETER LIBRARY for nuclear data evaluation.

The number of nuclear reaction channels open increases very rapidly with projectile energy, leading to the excitation of tens or even hundreds or residual nuclei. The tremendous effort needed for input preparation makes the evaluator community appreciate the strong necessity of having available, through direct code access, reliable standard parameter libraries.

The Nuclear Data Section of the IAEA (NDS), realizing the importance of this subject matter, has started a Coordinated Research Programme (CRP) aimed at the organization of a Reference Input Parameter Library (RIPL) for nuclear model codes. Accordingly, actions will be taken on a list of 7 groups of major input parameters:

- 1. atomic masses and related data,
- 2. discrete level schemes,
- average neutron resonance parameters,
- 4. optical model parameters,
- 5. level densities (total, partial, fission)
- 6. gamma-ray strength functions,
- 7. fission barriers.

Parameters 1,2,3 are model independent, whereas parameters 4, 5, 6, 7 are model dependent. In order to be able to decide which model parameters to be compiled, assessments achievable through critical intercomparisons of the physics underlying the different models must be made. This makes the two efforts being started by the NEANSC of OECD and by the NDS of IAEA, respectively, very much complementary, strictly interdependent, and mutually supportive. Therefore it will be one of the tasks of the s12 on NMV to provide theoretical support to the IAEA CRP on the RIPL.

2. NUCLEAR MODEL VALIDATION: EFFORT AND ORGANIZATION

2.1. The task.

The period of time in which s12 is foreseen to be active has been indicated as 3 years. It is impossible to imagine performing all the work in the physics involved up to a few hundred MeV in a 3 years time period. This period can be sufficient, however, to cast some light on the status of the art of all what may be needed in nuclear data evaluation in the energy range up to 200-300 MeV.

According to the indications of the NEANSC IEC, the NMV should consider all reaction mechanisms taking place below about one hundred MeV incident energy. One may suggest the pion production threshold as a more proper dividing line, in view of the implications of elementary particle production, a complication which requires different theoretical treatment. The energy domain so defined, typically involves reactions induced by light projectiles.

In terms of nuclear model approaches and theories currently used, the following scheme has been adopted:

SCHEMATIC REPRESENTATION OF CURRENTLY USED APPROACHES

	DISC	RETE	LEVELS	REG	ION	SHELL MC HARTREE- RPA		IBA	
NUCLEAR STRUCTURE	CONT	INUUM REGIO	.		TOTAL LVI	DNST	BCS FERMI SHELL		
					PARTIAL L	VI, DN	FERMI		
								LOCAL	
	OPT	MDT.	& DIR.	מאיץ פכווכו		SPHERICAL		NON LO	CAL
	V		w	tome s		COUPLED C	Η.		
	R-MA	TRIX	and R-N	ATR:	IX-S	SHELL MDL	APP. I	O LOW	MASSES
	THERM	MAL &	RESONA	ANCE	REG	SION			
	COMPO	OUND :	NUCLEUS	}		JSER-FESHB.			
NUCLEAR REACTIONS	PREEÇ	miri	BRIUM		HYB FKK WEI TAM	TITON MODE: RID MODEL: DENMULLER URA ET AL KA MODEL	ET AL		
	FISSI	ON M	ODELS		MAD	AMIC FISS: LAND MODE LIGER MODI	.₊		
				**************************************	G-A	BSORPTION		RESON	
	GAMMA	-RAY	REACT.		G-P	RODUCTION	DIREC COLLE PREEQ BREMS		IUM NG

2.2 The work organization.

Due to a complete lack of economical support the cooperation must take place mostly via e-mail, and this is additional

reason to split s12 into as small groups as possible.

Plenary or interdisciplinary sessions can be foreseen only at the occasion of IAEA meetings or international conferences as they arise. They will be good occasions for considering interdisciplinary aspects of links among different pieces of physics. At such occasions open problems will be raised and discussed.

From the schematic representation of table 1 one identify at least 8 different groups of specialists. The specialists on basic theory and on evaluation which have accepted to participate into the NMV activities are grouped here below, as agreed. A reference person (underlined) has been indicated for each group.

I. WORKING GROUP ON NUCLEAR STRUCTURE

S.	GRIMES (Oh	io Unive	ersity) U	SA
	HERMAN SATO	(guest (TCPK)	at ENEA) JAPAN	Italy
Y,	NAKAJIMA	(JAERI)		
	VENTURA	(ENEA)	Italy	
A.	V. IGNATYUK	(IPPE)	Russia	
	ZONGDI	(CNDC)	China	
C.	Y. FU	(ORNL)	USA	
G.	ROHR	(CBNM)	Belgium	

II. WORKING GROUP ON OPTICAL MODEL AND DIRECT REACTIONS

<u>F.</u>		DIETRICH (LLNL) USA
М. К. Y. Р.	MATOBA ISHIBASHI WATANABE G. YOUNG	(KYUSHU UN.) JAPAN (KYUSHU UN.) JAPAN (KYUSHU UN.) JAPAN (LANL) USA
V. B. V. D.	PRONYAEV STROHMAIER AVRIGEANU MADLAND BERSILLON	(IPPE) Russia (IRK) Austria (IPNE) Rumania (LANL) USA (CEA) France

III. WORKING GROUP ON R-MATRIX AND R-MATRIX-SHELL MODEL APPROACH FOR PARTICLE REACTIONS IN LIGHT ELEMENTS.

J.	HALE	(LANL)
D.	RESLER	(LLNL) USA
K.	SHIBATA	(JAERI) Japan
N.	LARSON	(ORNL) USA

IV. THERMAL AND RESONANCE REGION

F.	FROEHNER	(KFK) FRG
H.	DERRIEN	France

V. WORKING GROUP ON COMPOUND NUCLEUS AND PREEQUILIBRIUM REACTION MECHANISMS.

<u>M.</u>	<u>B.</u>	CHADWICK	(LLNL)	USA
p.	YOU	JNG	(LANL)	USA
			(LLNL)	
C.	Υ.	FU	(ORNL)	
ZHA	NG	JINGSHANG		
	CH3			Japan
Υ.	N.	SHUBIN	(IPPE)	Russia
				n Universitaet)
			FRG	, ,
Μ.	HEF	MAN	(quest	at ENEA) Italy
F.	DIE	ETRICH	(LLNL)	USA
A.	J.	KONING		
	KAI		. ,	(SITAET DRESDEN)
			FRG	
Υ.	WAT	'ANABE	(Kvushi	u Univ.) Japan
Α.	MAR	CINKOWSKI	(INR) F	Poland
		OHMAIER	(IRK) A	
•				

VI. WORKING GROUP ON FISSION MODELS

<u>.</u>	MADLAND	(LANL)	USA
	OHSAWA	(KINKI	UN.) JAPAN
	NG SHU NUAN	(CNDC)	China

VII. WORKING GROUP ON REACTION MECHANISMS INVOLVING GAMMA RAYS

<u>M.</u>	HERMAN	(guest at ENEA)	Italy
F.	DIETRICH	(LLNL) USA	
М.	IGASHIRA	(TOKYO INST.	OF
		TECHN.) JA	
Η.	KITAZAWA	(TOKYO INST.	OF
		TECHN.) JA	
E.	BETAK		
J.	KOPECKY	(ECN) Holland	•
Μ.	UHL	(IRK) Austria	

VIII. WORKING GROUP ON NUCLEAR MODEL INTERCOMPARISON

M.	BLANN	(LLNL) USA
P.	NAGEL	(NEA) OCDE
Р.	G. YOUNG	(LANL) USA
Μ.	B. CHADWICK	(LLNL) USA
F.	FABBRI	(ENEA) Italy
Μ.	HERMAN	(guest at ENEA) Italy
C.	Y. FU	(ÖRNL) USA
H.	KALKA	(UNIV. DRESDEN) FRG
S.	CHIBA	(JAERI) Japan
\mathtt{T} .	PUKAHORI	(JAERI) JAPAN
Μ.	UHL	(IRK) Austria
J.	A. KONING	(ECN) Holland
	N. SHUBIN	(IPPE) RUSSIA

B. STROHMAIER

(IRK) AUSTRIA

V. M. MASLOV

H. DERRIEN

(RPCPI) BELARUS

3. CODES RECOMMENDED FÖR CONSIDERATION

A list of codes has been suggested to consideration by the participants to the 1-st NMV meeting.

1. ALICE (BLANN)

Documented

2. GNASH (YOUNG)

Documented

3. GNASH-FKK (YOUNG-CHADWICK)

Documented

4. STAPREH93 (AVRIGEANU)

Documented

5. ALICE F (FUKAHORI)

6. SINCROS 2 (FUKAHORI-CHIBA)

7. CATHY (FUKAHORI)

Documented

8. ELIESE (FUKAHORI)

Documented

9. TNG (FU)

Documented

10. RHOTHERM (GRIMES)

SHELL MODEL + STATISTICAL

11. EDA (HALE)

GENERAL R-MATRIX, on Crav

12. ECIS 87 (PRONYAEV)

Multichannel coupling

13. ECIS 94 (KONING-RAYNAL)

Manual planned

14. EXIFON (KALKA)

Documented

15. EXIFON C (KALKA)

16. CAPSIES (KONING)

17. GLTLTLT (MADLAND)

Fission spectrum, average multeplicity, spectrum moment. NO DO-CUMENTATION.

18. RELOMP (MADLAND)

OPTICAL MODEL MEDIUM ENERGY SOME DOCUMENTATION

19. SNOOP 8 (MADLAND)

SHROEDINGER EO. OPTICAL MODEL +RELATIVISTIC ENERGY, 2-ND ORDER DIRAC, + STANDARD OPTICAL MODEL,

AS A PARTICULAR CASE

20. EMPIRE (MARCINLOWSKI)

HAUSER-FESHBACH + HYBRID. Documen-

ted.

21. EMPIRE (MARCINLOWSKI)

+ MSC FKK. Documented.

22. ANDRE' (MARCINLOWSKI)

FKK-MSD . Will be documented.

23. MISDO (ROHR)

Level Density Calculations.

24. STAPRE (UHL)

COMP. NUCL. + PREEQ. EXCITON MODEL, DOCUMENTED

25. FKK-GNASH (WATANABE)

26. SCIEDW (WATANABE)

DWBA

27. UNF (ZANG JINGSHANG)

EXCITON MDL, Documented.

28. FUNF (ZHANG JINGSHANG)

FISSILE NUCLEI

29. CUNF (ZHANG JINGSHANG)

CHARGED PARTICLE INDUCED REAC-

TIONS.

30. GUNF (ZHANG JINGSHANG)

GAMMA INDUCED REACTIONS.

31. IDA MODULAR SYSTEM (REFFO-FABERI-HERMAN)

AMLETO level density analysis
ESTIMA resonance analysis
ARGO reaction lists and Q-values
EURIALOgamma branchings
CERBERObinary cross sections
PENELOPEmultiple cascading emissions
NAUSICA as above + two gas preeq
CIRCE generalized optical model
EOLO valence capture
TELEMACO inclusive spectra calcts.
ULYSSES data processing.

- 33. PEGAS (BETAK-OBLOZINSKY)
- 34. GAMME (BONETTI-CHADWICK)
- 35. MAURINA (UHL)
- 36. GRAPE (GRUPPELAAR)

37. SAMMY (N. LARSON)

R-MATRIX

38. SCAT-2 (BERSILLON)

SPHERICAL OPTICAL MODEL

39. FLUKA (FERRARI et al.)

Documented, 1MeV-1000 GeV

4. OBJECTIVES AND MILESTONES

The working groups are invited to produce survey-documents with comments on the experience of the group members about models and tools they have produced and/or they have been using. The existing literature can be used, only it has to be reoriented according to the needs and scope of s12 activity. These documents will be collected by the coordinator and important aspects will be brought up and will constitute discussion elements until complete agreement is reached.

Previous code intercomparisons showed how the same code may yield user-dependent results. This experience puts in evidence the necessity of special attention to the methodological aspects of nuclear modelling. We should, therefore, like to promote any effort to have (in addition to code manuals) also theory and nuclear modelling guidelines. The analysis of previous code intercomparisons will offer useful indications about which types of interpretation ambiguities have given place to different results.

Our work will proceed accordingly, aiming to the following MAIN OBJECTIVES AND MILESTONES:

SURVEY OF THE DIFFERENT NUCLEAR MODELS DEVELOPED. (BY END OF '94).

implies a thorough critical analysis and intercomparison of the physics underlying the different approaches. To this end the work done by A.J. Koning will be helpful, see document NEA/NSC/DOC(92)12.

We will have to indicate what are the advantages and drawbacks of different model approximations and to give indications, as far as possible, of the reliability of results in terms of uncertainty figures.

We should provide guidance for the correct use of the models recommended, and give comprehensive indications about model parameters and inherent systematics.

INTERCOMPARISON OF MODEL CALCULATIONS CRITICAL AGAINST EXPERIMENT. (BY THE END OF '95).

Of necessity this step has to include:

- an assessment of the degree of confidence expected from the various approches which will be considered.
- critical intercomparison of the corresponding model b) We will be greatly helped by the code intercomparison

currently underway and coordinated by M. Blann and

- Similarly we will have to illustrate the advantages the range of validity of different codes considered for recommendation and release, along with user manuals.
- identification of critical experiments for theory testing and validation.
- identification of problems and gaps which may constitute d) the starting point of later efforts to be taken in the frame of international cooperation.

6. FUTURE METINGS

WE plan to have the 2-nd meeeting of s12 NMV in december 1994, by the time of the JEF-EFF meeting.

7. SHORT SUMMARY

++++ WHAT WE HAVE DONE AT THE FIRST \$12 NMV MEETING IS:

- # AGREE ON A WORK PROGRAMME
- # AGREE ON THE ACTIONS: WHO IS DOING WHAT
- # AGREE ON SOME MILESTONES

++++ WHAT IS NEEDED IS:

- # OUTLINE OF BASIC PHYSICS
- # VALIDITY LIMITS
- # SUCCESS AND FAILURES
- # DIFFICULTIES
- # RANGE OF APPLICABILITY IN E AND A
- # IS THERE ANY NEED OF EXPERIMENTAL BENCHMARKING
- # DIFFERENCES FROM OTHER APPROACHES
- # PARAMETERS AND PARAMETER SYST.
- # IF MORE APPROACHES ARE AVAILABLE SELECT A FEW AND SPECIFY THE REASONS OF THE CHOICE
- # PROVIDE REFERENCES FOR DETAILS
- # IS THE APPROACH CODED
- # IS THE CODE AVAILABLE
- # IS THE CODE PROVIDED WITH MANUAL
- # ARE MORE CODES AVAILABLE
- # IS A CODE INTERCOMP. NEEDED

++++ ACTIONS

1. Reffo: send out questionnaires on codes and models by july 15.

by Jury 15:

2. Grimes: will consider level density effects on cross sections.

3. All participants: write a note on the theory-approach of interest, respectively, by the end of 1994.

4. Madland: will include folding model-dispersive approaches to otpical model.

5. All participants: provide code manuals-documentation end of -94